

New Perspectives on the Treatment of Metastatic Renal Cell Carcinoma: An Introduction and Historical Overview

ROBERT J. MOTZER

Memorial Sloan-Kettering Cancer Center, New York, New York, USA

Key Words. Renal cell carcinoma • Kidney cancer • Metastatic • Vascular endothelial growth factor receptor • Mammalian target of rapamycin • mTOR

Disclosures: Robert J. Motzer: *Honoraria:* Eisai, Bristol-Myers Squibb; *Research funding/contracted research:* GlaxoSmithKline, Novartis, Pfizer.

The content of this article has been reviewed by independent peer reviewers to ensure that it is balanced, objective, and free from commercial bias. No financial relationships relevant to the content of this article have been disclosed by the independent peer reviewers.

In 2010, kidney or renal pelvis cancers are expected to be diagnosed in an estimated 58,240 individuals and to be responsible for 13,040 deaths in the U.S. [1]. Renal cell carcinoma (RCC) arises from the renal epithelium and represents approximately 85% of all kidney cancers [2]. Initial treatment for localized RCC is surgery, but 20%–30% of patients typically relapse, with a median time to relapse of 1–2 years [3]. In addition, approximately one third of patients with RCC present with metastatic disease [4]. Until recently, patients with RCC whose disease had metastasized had few treatment options. Cytokines (interleukin-2 and interferon- α) were the standard therapy but were associated with relatively low response rates and significant toxicity [5–7].

The treatment of metastatic RCC has changed dramatically in the past 5 years with the availability of new treatment options (Fig. 1). Molecular research into the pathogenesis of RCC has provided valuable information on the signaling pathways known to be altered in RCC, including vascular endothelial growth factor (VEGF) and its receptor (VEGFR) and the mammalian target of rapamycin (mTOR), a key kinase regulating cell growth and prolifer-

ation, cellular metabolism, and angiogenesis. Molecularly targeted therapies were developed specifically to target these signal transduction pathways. These include bevacizumab, a monoclonal antibody targeting VEGF; the tyrosine kinase inhibitors sunitinib, sorafenib, and pazopanib, which target VEGFR and other receptor tyrosine kinases; and everolimus and temsirolimus, selective mTOR inhibitors. In pivotal phase III trials, bevacizumab, sunitinib, sorafenib, pazopanib, and temsirolimus have shown clinical benefit for patients with treatment-naïve or cytokine-pre-treated RCC by prolonging progression-free or overall survival (temsirolimus only) times [8–15]. Clinical benefit was observed in patients with metastatic RCC treated with everolimus as a second-line therapy after failure of sunitinib or sorafenib, as demonstrated by the more than double median progression-free survival time, compared with placebo [16, 17].

Because of their distinct molecular targets, the safety profiles of these agents are drug class-specific. The majority of adverse events observed in phase III trials of these agents were grade 1 or 2, although grade 3 or 4 toxicities were observed in 10%–20% of patients [18]. Consequently,

Correspondence: Robert J. Motzer, M.D., Memorial Sloan-Kettering Cancer Center, 1275 York Avenue, New York, New York 10021, USA. Telephone: 646-422-4312; Fax: 212-988-0719; e-mail: motzerr@mskcc.org Received September 9, 2010; accepted for publication December 1, 2010. ©AlphaMed Press 1083-7159/2011/\$30.00/0 doi: 10.1634/theoncologist.2011-S2-01




Figure 1. Timeline of treatments for renal cell carcinoma.

Abbreviations: EMA, European Medicines Agency; FDA, U.S. Food and Drug Administration; IFN, interferon; IL, interleukin.

patients treated with targeted therapies require close monitoring with specific management algorithms developed to optimize patient benefit. Given the generally poor prognosis of patients with metastatic RCC and the toxicity associated with therapy, health-related quality of life and patient-reported outcomes have become an important consideration in this patient population because survival for these patients is prolonged [19].

The efficacy and safety profiles of these targeted agents have resulted in U.S. and European regulatory approvals and updated recommendations within the U.S. (National Comprehensive Cancer Network) and European (European Organisation for Research and Treatment of Cancer, European Society of Medical Oncology, European Association of Urology) clinical practice guidelines, thus revolutionizing the treatment of patients with metastatic RCC [3, 20–22]. Given the availability of multiple new targeted agents within a short period of time, the optimal sequence and/or combination of therapies for patients with RCC is an increasingly active area of research. Ongoing clinical trials currently are investigating combination and sequential ther-

apy with these targeted agents to identify regimens that will benefit patients with advanced RCC.

In this supplement, we review the pathogenesis and diagnosis of RCC and the efficacy and safety of currently available targeted therapies, including recommendations for adverse event management. We also discuss quality of life considerations when selecting targeted therapies for RCC and current treatment guidelines based on evidence-based medicine. In addition, preliminary data comparing the cost-effectiveness of these therapies in the first- and second-line settings are discussed. Finally, we highlight future directions including trials to determine the optimal setting and sequence of these agents and those that investigate novel agents, combinations, or sequential therapies to improve patients' survival outcomes.

ACKNOWLEDGMENTS

The author takes full responsibility for the content of the paper but thanks Victoria Robb, Ph.D., and Amy Zannikos, Pharm.D., supported by Novartis Pharmaceuticals Corporation, for their assistance in manuscript writing and editing.

REFERENCES

- 1 American Cancer Society. Cancer Facts & Figures 2010. Atlanta: American Cancer Society; 2010. Available at <http://www.cancer.org/acs/groups/content/@nho/documents/document/acspc-024113.pdf>, accessed August 25, 2010.
- 2 Cohen HT, McGovern FJ. Renal-cell carcinoma. *N Engl J Med* 2005;353:2477–2490.
- 3 NCCN Clinical Practice Guidelines in Oncology™ Kidney Cancer. Version 2.2010. ©2009 National Comprehensive Cancer Network, Inc. Avail-

- able at http://www.nccn.org/professionals/physician_gls/f_guidelines.asp#site, accessed July 30, 2010.
- 4 Altekruze SF, Kosary CL, Krapcho M et al. eds. SEER Stat Fact Sheets: Kidney and Renal Pelvis. SEER Cancer Statistics Review, 1975–2007. Bethesda, MD: National Cancer Institute. Available at http://seer.cancer.gov/csr/1975_2007, accessed July 30, 2010.
- 5 Belldegrun AS, Klatte T, Shuch B et al. Cancer-specific survival outcomes among patients treated during the cytokine era of kidney cancer (1989–2005): A benchmark for emerging targeted cancer therapies. *Cancer* 2008; 113:2457–2463.
- 6 Oudard S, George D, Medioni J et al. Treatment options in renal cell carcinoma: past, present and future. *Ann Oncol* 2007;18(suppl 10):x25–x31.
- 7 Atkins MB, Regan M, McDermott D. Update on the role of interleukin 2 and other cytokines in the treatment of patients with stage IV renal carcinoma. *Clin Cancer Res* 2004;10:6342s–6346s.
- 8 Escudier B, Pluzanska A, Koralewski P et al. Bevacizumab plus interferon alfa-2a for treatment of metastatic renal cell carcinoma: A randomised, double-blind phase III trial. *Lancet* 2007;370:2103–2111.
- 9 Escudier B, Eisen T, Stadler WM et al. Sorafenib in advanced clear-cell renal-cell carcinoma. *N Engl J Med* 2007;356:125–134.
- 10 Escudier B, Eisen T, Stadler WM et al. Sorafenib for treatment of renal cell carcinoma: Final efficacy and safety results of the phase III treatment approaches in renal cancer global evaluation trial. *J Clin Oncol* 2009;27:3312–3318.
- 11 Motzer RJ, Hutson TE, Tomczak P et al. Overall survival and updated results for sunitinib compared with interferon alfa in patients with metastatic renal cell carcinoma. *J Clin Oncol* 2009;27:3584–3590.
- 12 Escudier B, Bellmunt J, Négrier S et al. Phase III trial of bevacizumab plus interferon alfa-2a in patients with metastatic renal cell carcinoma (AVOREN): Final analysis of overall survival. *J Clin Oncol* 2010;28:2144–2150.
- 13 Motzer RJ, Hutson TE, Tomczak P et al. Sunitinib versus interferon alfa in metastatic renal-cell carcinoma. *N Engl J Med* 2007;356:115–124.
- 14 Sternberg CN, Davis ID, Mardiak J et al. Pazopanib in locally advanced or metastatic renal cell carcinoma: Results of a randomized phase III trial. *J Clin Oncol* 2010;28:1061–1068.
- 15 Hudes G, Carducci M, Tomczak P et al. Temsirolimus, interferon alfa, or both for advanced renal-cell carcinoma. *N Engl J Med* 2007;356:2271–2281.
- 16 Motzer RJ, Escudier B, Oudard S et al. Efficacy of everolimus in advanced renal cell carcinoma: A double-blind, randomised, placebo-controlled phase III trial. *Lancet* 2008;372:449–456.
- 17 Motzer RJ, Escudier B, Oudard S et al. Phase 3 trial of everolimus for metastatic renal cell carcinoma: Final results and analysis of prognostic factors. *Cancer* 2010;116:4256–4265.
- 18 Shepard DR, Garcia JA. Toxicity associated with the long-term use of targeted therapies in patients with advanced renal cell carcinoma. *Expert Rev Anticancer Ther* 2009;9:795–805.
- 19 Cella D. Quality of life in patients with metastatic renal cell carcinoma: The importance of patient-reported outcomes. *Cancer Treat Rev* 2009;35:733–737.
- 20 EAU Guidelines on Renal Cell Carcinoma. ©2009 European Association of Urology. Available at <http://www.uroweb.org>, accessed July 30, 2010.
- 21 de Reijke TM, Bellmunt J, van Poppel H et al. EORTC-GU group expert opinion on metastatic renal cell cancer. *Eur J Cancer* 2009;45:765–773.
- 22 Escudier B, Katja V; ESMO Guidelines Working Group. Renal cell carcinoma: ESMO clinical recommendations for diagnosis, treatment and follow-up. *Ann Oncol* 2009;20(suppl 4):81–82.